

Scope of Practice

Approved by

State Board of Emergency Medical, Fire and Transportation Services and the Ohio Department of Public Safety, Division of EMS

This document offers an "at-a-glance" view of the Scope of Practice for Emergency Medical Responders (EMR), Emergency Medical Technicians (EMT), Advanced Emergency Medical Technicians (AEMT), and Paramedics as approved by the State Board of Emergency Medical, Fire, and Transportation Services (EMFTS Board). The authorized services can be found in sections 4765.35 (FR/EMR), 4765.37 (EMT-B/EMT), 4765.38 (EMT- I/AEMT), and 4765.39 (EMT-P/Paramedic) of the Revised Code. The scopes of practice can be found in rules 4765-12-04 (EMR), 4765-15-04 (EMT), 4765-16-04 (AEMT), and 4765-17-03 (Paramedic) of the Administrative Code. The Ohio EMS scope of practice authorized by the EMFTS Board applies to emergency and non-emergency settings.

Performance of services outlined in this document and in the aforementioned code sections, shall only be performed if the EMR, EMT, AEMT, and Paramedic have received training as part of an initial certification course or through subsequent training approved by the EMFTS Board. If specific training has not been specified by the EMFTS Board, the EMR, EMT, AEMT, and Paramedic must have received training regarding such services approved by the local medical director before performing those services.

The individual medical director of each EMS agency may limit or ask that providers obtain approval from medical direction for certain treatments. Each medical director may need to tailor and revise their protocol to meet their community's needs and to fit their region and individual practice, but they must ensure that all protocols remain within the approved scope of practice. EMS medical directors are reminded that they are not permitted to expand or exceed the scope of practice for EMS providers which has been authorized by the EMFTS Board; however, they may provide clarifications or limitations on services that are permitted.

EMS medical directors and EMS providers are strongly encouraged to review the EMFTS Board's policy statement "Regarding EMS Provider Prehospital Transport of Patients with Pre-Existing Medical Devices or Drug Administrations" dated February 2018 (attached at the end of this document). This statement clarifies how EMS providers, in the prehospital setting, should deal with medical devices and medicine administrations that are outside their scope of practice.

Pursuant to rule 4765-6-04 of the Administrative Code, the EMFTS Board may allow EMRs, EMTs, AEMTs, and Paramedics to perform services beyond their respective scopes of practices as part of a board-approved research study. An entity must submit a research proposal to the EMFTS Board in accordance with the requirements of rule 4765-6-04 of the Administrative Code. The EMFTS Board is not obligated to approve the proposed research study nor accept any recommendation to permanently amend the scope of practice.

Updated 11/19/03; 5/17/05; 10/26/05; 10/17/07; 3/8/12; 8/22/13, 10/16/13, 12/18/13, 4/16/2014, 10/19/16, 2/15/17, 10/18/17, 6/20/18, 4/1/20



State Board Emergency Medical, Fire and Transportation Services Ohio Department of Public Safety, Division of EMS

Updated April 1, 2020

	Airway Management	EMR	EMT	AEMT	PARAMEDIC
1	Open and maintain the airway	Х	Х	X	X
2	Oropharyngeal airway adjunct	Х	Х	Х	Х
3	Nasopharyngeal airway adjunct	Х	Х	X	X
4	Manual removal of obstructed airway	Х	X	X	X
5	Laryngoscopy for removal of airway obstruction			Х	х
6	Oral suctioning	X	Х	х	Х
7	Endotracheal (ET) tube suctioning through a previously established airway or a stoma		х	х	х
8	Tracheostomy tube replacement			Х	X
9	Cricothyrotomy, surgical				х
10	Cricothyrotomy, needle				Х
11	Apply and obtain readings of pulse oximeter, CO-oximeter, and capnography or capnometry equipment	х	х	х	х
12	Oxygen administration				
	a. Nasal cannula	Х	Х	Х	X
	b. Non-rebreather mask	х	х	х	х
	c. Mouth-to-barrier devices	Х	Х	х	х
	d. Partial rebreather mask		Х	Х	Х
	e. Venturi mask		Х	х	х
13	Ventilation management				
	a. Bag valve mask	Х	Х	х	Х
	b. Ventilation with a flow-restricted oxygen-powered device	Х	х	х	х
	c. Positive pressure ventilation devices (manually triggered or automatic ventilators)		х	х	x
14	Ventilator management - 16 years of age or older				х

	Airway Management (cont'd)	EMR	EMT	AEMT	PARAMEDIC
15	Non-emergent ambulance transport of a stable patient less than 16 years of age who has a chronic condition requiring a tracheostomy tube and a ventilator provided the patient's caregiver accompanies the patient during transport. The caregiver must have received appropriate training in use of the patient's ventilator. A caregiver is not required to accompany the patient if the patient is accompanied by an Ohio licensed registered nurse or respiratory therapist, or other appropriately trained and licensed Ohio healthcare provider.				X
16	Orotracheal intubation				Х
	a. Apneic patients			Х	Х
	b. Pulseless <u>and</u> apneic patients			Х	Х
17	Nasotracheal intubation				Х
18	Dual lumen airway				Х
	a. Apneic patients			Х	Х
	b. Pulseless <u>and</u> apneic patients		х	Х	Х
19	Extraglottic airways				Х
	a. Apneic patients			Х	Х
	b. Pulseless <u>and</u> apneic patients		Х	Х	Х
20	CPAP administration and management		х	Х	Х
21	BiPAP administration and management				Х
22	Positive end-expiratory pressure (PEEP)				Х
23	End tidal CO₂ monitoring and detecting		Х	Х	Х
24	Oxygen humidifier equipment application and monitoring		Х	Х	Х
25	Chest tube monitoring and management				Х
26	Nasogastric (NG) tube placement				Х
27	Orogastric (OG) tube placement				Х

	Cardiac Management	EMR	EMT	AEMT	PARAMEDIC
1	Cardiopulmonary resuscitation (CPR)	х	X	X	X
2	Chest compression assist devices	Х	х	x	X
3	Automated external defibrillator (use of an AED)	х	х	x	х
4	Manual defibrillation			х	Х

	Cardiac Management (Cont'd)	EMR	EMT	AEMT	PARAMEDIC
5	Negative impedance threshold devices		х	х	х
6	Administration of cardiac medication				Х
7	Set up cardiac monitor in the presence of an AEMT or Paramedic		х		
8	Cardiac monitor strip interpretation			Х	Х
9	Cardioversion				Х
10	Carotid massage				Х
11	Transcutaneous cardiac pacing				Х
12	12-lead EKG performance and interpretation				Х
	12-lead EKG application assisting a				
13	Paramedic who is present		Х	X	
14	12-lead EKG set up and application for electronic transmission ^A		x	x	х

^A An EMT or AEMT may set up and apply a 12-lead electrocardiogram when assisting a Paramedic or for the purposes of electronic transmission <u>if all of the following conditions are met</u>: 1) performed in accordance with written protocol; 2) EMT or AEMT shall not interpret the electrocardiogram; 3) delay in patient transport is minimized; and 4) EKG is used in conjunction with destination protocols approved by the local medical director.

	Medical Management	EMR	EMT	AEMT	PARAMEDIC
1	Epinephrine administration via auto- injector	X	х	х	Х
2	Epinephrine administration via SQ or IM routes			x	Х
3	Epinephrine administration via IV or IO route				Х
4	Aspirin administration		Х	х	Х
5	Oral glucose administration		X	х	Х
6	Activated charcoal administration		X	х	Х
7	Nitroglycerin administration (patient assisted) ^B		х	x	Х
8	Nitroglycerin administration (non- patient assisted)			х	Х
9	Aerosolized or nebulized medications administration (patient assisted) ^B		х	х	х
10	Administration of aerosolized or nebulized medications (non-patient assisted)			х	х
11	Naloxone administration via auto- injector	Х	х	x	Х
12	Naloxone administration via intranasal route	Х	х	х	Х
13	Naloxone administration via ETT, IM, IV, IO, or SQ routes			х	Х
14	Medication administration (protocolapproved) ^c			х	х

	Medical Management (Cont'd)	EMR	EMT	AEMT	PARAMEDIC
15	Administration of intranasal medications (in addition to naloxone) ^c			х	х
16	Immunizations for influenza to firefighters, EMTs, AEMTs, or Paramedics (ORC 4765.391)				х
17	Set up of IV administration kit in the presence of an AEMT or Paramedic		х		
18	Transport of central/peripheral IV without an infusion		х	х	х
19	IV maintenance and fluid administration			х	Х
20	Maintenance of medicated IV fluids				Х
21	Central line monitoring				Х
22	IV infusion pump				Х
23	Intraosseous needle insertion			х	Х
24	Saline lock initiation			х	Х
25	Peripheral IV blood specimens			х	Х
26	Maintenance of blood administration				Х
27	Thrombolytic therapy initiation and monitoring				Х

^B Patient Assisted Definition: May assist with 1) patient's prescription upon patient request and with written protocol – OR – 2) EMS-provided medications with verbal medical direction.

^c See "AEMT Medications Approved by the EMFTS Board."

	Trauma Management	EMR	EMT	AEMT	PARAMEDIC
1	PASG		х	х	х
2	Long spine board	х	х	х	Х
3	Short spine board	х	х	х	Х
4	Splinting devices	х	х	х	Х
5	Traction splint		х	х	Х
6	Cervical immobilization device (CID)	х	х	х	х
7	Helmet removal		х	х	Х
8	Rapid extrication procedures		х	х	х
9	Needle decompression of the chest			х	х
10	Soft tissue management	х	х	х	х
11	Management of suspected fractures	х	х	х	х
12	Controlling of hemorrhage	х	х	х	х

	Basic Performances	EMR	EMT	AEMT	PARAMEDIC
1	Body substance isolation precaution/administration	х	Х	Х	х
2	Taking and recording of vital signs	Х	х	Х	х

	Basic Performances	EMR	EMT	AEMT	PARAMEDIC
3	Patient Care Report (PCR) documentation	х	х	Х	Х
4	Trauma triage determination per OAC 4765-14-02	Х	X	x	Х

	Additional Services	EMR	EMT	AEMT	PARAMEDIC
1	Emergency childbirth management P	x	x	x	Х
2	Glucose monitoring system use (with				
	Clinical Laboratory Improvement				
	Amendments (CLIA) waiver in place		X	X	X
3	Blood analysis				Х
4	Eye irrigation	Х	Х	Х	Х
5	Eye irrigation with Morgan lens				X
6	Maintenance of blood administration				X
7	Thrombolytic therapy initiation and monitoring				Х
[₽] An El	MR may only assist with emergency childbi	rth manageme	nt.		

Emergency Medical Services in Hospital	EM R	EMT	AEMT	PARAMEDIC
In a hospital, an EMT, AEMT or Paramedic may perform emergency medical services in accordance with the following conditions: only in the hospital's emergency department (ED) or while moving a patient between the ED and another part of the hospital; only under the direction and supervision of a physician, a physician assistant designated by a physician, or a RN designated by a physician (ORC 4765.36). The EMT, AEMT, or Paramedic cannot perform any service outside the scope of practice of his or her certificate to practice.		х	х	х
Additional Services in a Declared	EM	EMT	AEMT	PARAMEDIC
In the event of an emergency declared by the governor that affects the public's health, an EMS provider may perform immunizations and administer drugs or dangerous drugs, in relation to the emergency, provided the EMS provider is under physician medical direction and has received appropriate training regarding the administration of such immunizations and/or drugs. (OAC 4765-6-03)	X	х	х	х

Nerve Agent or Organophosphate Release	EM R	EMT	AEMT	PARAMEDIC
An EMS provider may administer drugs or dangerous drugs contained within a nerve agent antidote auto-injector kit, including a MARK I* kit, in response to suspected or known exposure to a nerve or organophosphate agent provided the EMS provider is under physician medical direction and has received appropriate training regarding the administration of such drugs within the nerve agent antidote auto-injector kit. (OAC 4765-6-05)	х	х	X	x
Withdrawing of Blood for Evidence	EMR	EMT	AEMT	PARAMEDIC
Collection				
Withdraw blood for the purpose of determining the alcohol, drug, controlled substance, metabolite of a controlled substance, or combination content of the whole blood, blood serum, or blood plasma only if the medical director provides authorization, a written protocol, and training. It may only be performed in the course of the provision of emergency medical treatment and at the request of a law enforcement officer, and only in response to a request for emergency medical treatment and transport to a health care facility. A clinically competent patient may refuse transport. Withdrawal of blood shall not be done: 1. If the physical welfare of the patient, EMS provider, or other person would be endangered 2. If it causes an unreasonable delay in treatment or transport of the patient or any other person 3. Consent of the patient is not obtained (an unconscious person or a person with a condition rendering the person incapable of refusal shall be deemed to have consented) 4. From a pre-existing central venous access device 5. Withdrawal of blood violates any rule in this chapter (OAC 4765-6) 6. The person is deceased (OAC 4765-6-06)			X	x


AEMT Medication Administration Approved by the EMFTS Board

A certified AEMT may administer medications from the following list, provided the AEMT is under physician medical direction and has received appropriate training regarding the administration of such medications. A medication that does not appear on the following list SHALL NOT be added to the department's AEMT protocol.

Lidocaine for pain relief after intraosseous needle insertions
Nalbuphine
Naloxone
Narcotics or other analgesics for pain relief
Nitrous oxide
Oral ondansetron ^E
Sublingual nitroglycerin

^EA certified AEMT may administer oral ondansetron for patients age 12 years or older. .

The approved route of administration of any specific medication is stated in the respective EMT, AEMT, and Paramedic curriculum. The EMS provider shall administer medications only via the route addressed in each respective curriculum and consistent with their level of training.



1970 West Broad Street, P.O. Box 182073, Columbus, Ohio 43218-2073

(614) 466-9447 / (614) 466-9461 FAX

Thomas Allenstein, Chair

The State Board of Emergency Medical, Fire, and Transportation Services ("EMFTS Board") issues the following statement:

> Regarding EMS Provider Prehospital Transport of Patients with Pre-Existing Medical Devices or Drug Administrations February 2018

This statement is an attempt to provide general information about the above issue facing EMS providers. It should not be treated as legal advice or medical direction. For direct advice regarding a particular scenario, please consult with your medical director and legal counsel. Although the following statement represents the EMS Board's general position on the above issue, this statement in no way precludes the EMFTS Board from taking disciplinary action in a particular case if necessary. Any potential complaints brought before the EMFTS Board will be decided on a case-bycase basis.

Introduction:

The EMFTS Board and the Ohio Department of Public Safety, Division of Emergency Medical Services, has developed a defined scope of practice for EMS providers. It is maintained in matrix form and available on-line as a reference for public access. This scope of practice addresses all levels of EMS providers and has been approved by the EMFTS Board. Updates to the scope of practice are made as necessary and after approval by the EMFTS Board.

From time to time, EMS providers are confronted on-scene with patients with preexisting medical situations not included or addressed in their respective EMFTS Board approved scope of practice. Specifically, patients with pre-existing medical devices and drug administrations requiring prehospital EMS service are becoming more commonplace. The intent of this position paper is to address the EMS provider's approach to that prehospital patient with a pre-existing physician-ordered medical device or drug administration ("MDDA") not covered in the provider's scope of practice.

Discussion:

In general, the EMS provider should maintain the pre-existing MDDA and transport the patient to the appropriate facility. There is no expectation that the EMS provider will initiate, adjust, or discontinue the preexisting MDDA. This implies that the EMS provider will maintain and continue care so that the patient can be transported.

The EMS provider is expected to follow local protocols regarding the overall evaluation, treatment, and transportation of this type of prehospital patient requiring EMS service. It applies to EMS provider situations where alternative transportation and care is not available or practical (prehospital or "911 scene response"). It implies that the most appropriate and available level of EMS provider will respond to the request for prehospital EMS service. It also implies that the patient requires the pre-existing MDDA and it is not feasible or appropriate to transport the patient without the pre-existing MDDA.

The number and type of pre-existing MDDAs currently or potentially encountered by the EMS provider in the community setting is extensive and may change frequently. The intent of this position paper is not to provide an inclusive list of pre-existing MDDAs. However, as a guideline for the EMS provider, current pre-existing MDDAs may include ventilatory adjuncts (CPAP, BiPAP), continuous or intermittent IV medication infusions

State Board of Emergency Medical, Fire, and Transportation Services Page 2

(analgesics, antibiotics, chemotherapeutic agents, vasopressors, cardiac drugs), continuous gastric or parenteral infusion of nutrition, and nontraditional out-of-hospital drug infusion routes (subcutaneous infusaports, central venous access lines, direct subcutaneous infusions, self-contained implanted pumps).

Conclusion:

In conclusion, the EMS provider confronted with a prehospital patient with a pre-existing physician-ordered medical device or drug administration not covered in the EMS provider's respective scope of practice should provide usual care and transportation while maintaining the pre-existing MDDA, if applicable. Concerns or questions regarding real-time events associated with a pre-existing MDDA should be directed to the relevant physician providing medical direction. Concerns or questions regarding previous, recurrent, or future prehospital transportations with a pre-existing MDDA should be directed to the appropriate EMS medical director and legal counsel.